This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| (51) International Patent Classification 5: | , | (11) International Publication Number: WO 91/18588 |
|---|--|--|
| A61K 7/48, 7/075, 7/50 A61K 7/00 | A1 | (43) International Publication Date: 12 December 1991 (12.12.91) |
| (21) International Application Number: PCT/FI (22) International Filing Date: 5 June 1991 (30) Priority data: 533,684 5 June 1990 (05.06.90) (71) Applicant: CULTOR LTD. [FI/FI]; P.O. Box 1 kinportti 2, SF-00241 Helsinki (FI). (72) Inventor: JUTILA, Kirsti, Maarit; Rantapolku 2 02170 Espoo (FI). (74) Agent: RINKINEN, Raili; Cultor Ltd., P.O. Box likinportti 2, SF-00241 Helsinki (FI). | (05.06.9 05, Kyl B 10, S 105, K | patent), CH (European patent), DE, DÉ (European patent), DK (European patent), ES (European patent), FR (European patent), GB, GB (European patent), GR (European patent), IT (European patent), LU (European patent), NL (European patent), SE (European patent). Published With international search report. |
| (54) Title: A METHOD OF REDUCING THE IRRI | TATIN | IG PROPERTIES OF A COSMETIC COMPOSITION |

(57) Abstract

A method for substantially reducing skin irritation and improving moisturizing properties of cosmetics by incorporating a trimethylglycine compound in an effective amount, and cosmetic products produced thereby are disclosed.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| AT AU BB BE BP BG BJ BR CA CF CG CH CI CM CS DE DK | Austria Australia Barbados Belgium Burkina Faso Bulgaria Benin Brazil Canada Central African Republic Congo Switzerland Côte d'Ivoire Czechoslovakia Germany Denmark | ES PI FR GA GB GN HU IT JP KP KR LI LK LU MC | Spain Finland France Oabon United Kingdom Guinea Greece Hungary Italy Japan Democratic People's Republic of Korea Republic of Korea Liechtenstein Sri Lanka Luxembourg Monaco | MG ML MN MW NL NO PL SD SE SN SU TD TG US | Madagascar Mali Mongolia Mauritania Malawi Netherlands Norway Poland Romania Sudan Sweden Senegal Soviet Union Chad Togo United States of America |
|--|--|--|---|---|---|
|--|--|--|---|---|---|

3.

WO 91/18588

1

A METHOD OF REDUCING THE IRRITATING PROPERTIES OF A COSMETIC COMPOSITION

BACKGROUND OF THE INVENTION

5

10

15

20

25

30

35

There is an increasing demand for better and more efficient raw materials for cosmetics and related products. Even the most efficient raw materials can be very irritating or sensitizing for human skin, thus preventing their regular use.

Some of the active compounds in creams, decorative cosmetics, etc., may cause irritation or skin-sensitization. Such active ingredients may include emollients, barrier agents, healing agents, humectants, emulsifiers, preservatives, perfume oils, coloring agents, and/or surfactants, particularly anionic surfactants (e.g., sodium lauryl ether sulphates), and several compounds for skin cleansing (which contain kerosenses, solvents and disinfectants in addition to hard surfactants). Other cosmetics such as deodorants containing ethanol are recognized as having potentially irritating or sensitizing properties. In addition, this problem is often found with many other medicated products.

Cationic compounds such as cetyltrimethylammonium chloride, alkyldimethylbenzylammonium, alkylisoquinidinium, and alkylpyridinium halides are commonly
used as hair conditioners. In small amounts, these
quaternary ammonium derivatives improve hair manageability.

One drawback of including quaternary ammonium compounds in cosmetics such as shampoos, hair conditioners, creams, lotions, etc. is the fact that great care must be taken when introducing them into commercial products, as most of these materials are known to be skin irritants. More particularly, it is known from the literature that the quaternary compounds of synthetic

origin, particularly at higher concentrations (over 10 percent), cause skin irritation and cauterization.

This problem is overcome in hair conditioners and hair rinses because these preparations are diluted about 15 times with water to form the actual rinse used on the hair. Thus, in general, it has been found that when quaternary compounds are present in concentrations upon dilution with water of 0.1 percent, they are usually still safe and effective on the hair shaft.

Certain other cationic compounds introduced into conditioners have been touted as having a lower level of irritancy than earlier compounds. Examples of such include ethoxylated quaternary ammonium phosphates, quaternized fatty acid amino-amides derived from lanolic 15 acids and mink oil, and N-acyl colaminoformylmethylpyridinium derivatives.

Quaternary ammonium salts have also previously been used at concentrations below 0.1 percent as antiseptics in preparations such as aftershave lotions and 20 baby lotions and creams.

One hair treatment composition, described in U.S. Patent No. 4,752,467 (Konrad et al.), contains 0.1-25 percent by weight betaine and 0.1-10 percent by weight of an aliphatic organic acid such as citric acid. 25 The combination of the betaine/aliphatic organic acid is said to provide synergistic conditioning properties which improve the condition of the hair structure, and act as an antioxidant and as a buffer.

DE-Patent No. 1911144 (Medisan Ab) describes a 30 skin-treating composition for treating abnormal skin conditions or as a vehicle for other medicaments, comprising an aqueous phase in which urea and lactic acid are dissolved. The mode of operation of this composition 35 _relates to the skin-softening properties of urea and the keratolytic effect of lactic acid. Optional components

of this composition include emulsifiers and amino acid derivatives such as betaine.

Other cosmetic formulations also include betai-Betaine is a general name for an organic tertiary amine with three free organic radicals. Known formulations include betaines in which at least one of the free radicals is a long fatty acid derivative based radical. U.S. Patent No. 4,490,355 (Desai) describes a mixture of cocoamidopropyl betaine and oleomidopropyl betaine which when included in cosmetics is said to improve thickening and foam boosting properties. U.S. Patent No. 4,654,161 (Kollmeier et al.) describes the use of organopolysilaxanes that have betaine groups in hair cosmetics. use of these siloxane derivatives is said to provide improved compatibility with anionic additives (surfactants) and less irritation than prior organopolysiloxanes with quaternary ammonium groups. EP-Patent No. 286261 (Redken Laboratories) describes the use of zwitterions such as taurines and betaines to increase the 20 substantivity of hydrolysed proteins in keratinous tissues.

It is an object of the present invention to provide a method of reducing the skin irritating properties of a cosmetic composition containing substantially no hydrolysed protein.

It is another object of this invention to provide a cosmetic composition which has improved skin compatibility and moisturizing properties.

30 SUMMARY OF THE INVENTION

It has now been surprisingly discovered that adding an effective amount of a quaternary ammonium like compound which does not have any skin irritating properties to cosmetic compositions reduces the skin irritating 35 properties of skin irritating cosmetics and improves their skin compatibility and moisturizing properties. The present invention also relates to the method of

-

10

15

20

25

reducing the skin irritating properties of a cosmetic composition.

The quaternary ammonium compound, preferably obtained from natural sources, is methanaminium, 1-carboxy-N,N,N-trimethyl-,hydroxide inner salt monohydrate or the corresponding anhydride thereof (also referred to as trimethylglycine or trimethylglycine anhydride respectively).

The present invention also relates to a cosmetic composition comprising one or more cosmetic ingredients selected from the group consisting of an active agent, an emollient, a barrier agent, a healing agent, a humectant, an emulsifier, a preservative, a perfume oil, a coloring agent, a deodorizing agent, a surfactant, a solvent and mixtures of the above and one or more of the cosmetic ingredient having irritating properties when applied to human skin, and trimethylglycine or anhydride in an amount effective to substantially neutralize the skin-irritating properties of the cosmetic ingredient.

The dosage of trimethylglycine (or its anhydride) is from about 0.1 to about 50, preferably 1 - 15, percent by weight.

The cosmetic compositions of the present invention include skin cleansers, body shampoos, hair conditioners, skin creams or lotions, deodorants, etc.

DETAILED DESCRIPTION

The addition of trimethylglycine, a natural quaternary ammonium like compound methanaminium,

1-carboxy-N,N,N-trimethyl-,hydroxide inner salt monohydrate or the corresponding anhydride (trimethylglycine/ah)

÷.

WO 91/18588 PCT/FI91/00170

5

has now surprisingly been found to reduce the skin irritation properties or sensitisation reactions of cosmetic compounds including irritating ingredients such as sodium lauryl ether sulphate, and other anionic surfactants, cetyltrimethylammonium chloride, alkyldimethylbenzylammonium, and alkylisoquinidinium, alkylpyridinium halides, solvents etc. Trimethylglycine (or anhydride thereof) in question has been found to improve the skin compatibility and moisturizing characteristics of cosmetic compositions. Problems mentioned above are often found in products containing irritants, including skin and hand cleaners, skin lotions, skin creams, as well as medicated products. The dosage of methanaminium, 1-carboxy-N,N,N-trimethyl-, hydroxide, inner salt monohydrate or corresponding

The invention is further elucidated by the following examples, which are not meant to limit the claims.

anhydride is between 0,1 and 50%, preferably 1 - 15 w/w.

20

Examples 1-16

The effect of methanaminium, 1-carboxy-N,N,N-trimethyl,-hydroxide, inner salt monohydrate on skin compatibility and irritation were tested by human patch testing.

The patch testing was conducted as follows. In each of these Examples, the test materials were applied on the upper outer arm and the application arm was occluded with a patch for an initial contact period of 24 hours. The patches were then removed and the skin reaction was assessed using a numerical scoring system one hour later.

In order to assess the degree of irritancy, several signs symptomatic of skin irritation and their severity were noted using the scoring system below:

.

Service of the specific of the service of

| | | | . 8 |
|---|------------------|-----------|---------------------------------------|
| | 0. | = | DO vicina |
| | 1 | = | no visible relevant reaction |
| | 2 | = | reaction just present slight reaction |
| 5 | 3 | = | moderate reaction |
| | Mh a sa ta | = | |
| | The signs of irr | itation - | reaction |

The signs of irritation were awarded different ratings of significance as follows:

| | - an rollows: | | |
|----|-------------------------|-------------------|--------|
| 10 | <u>SIGN</u> Vesicles | ABBREVIATION V | RATING |
| 10 | Odema | v Oe | 5 |
| | Erythema | R | 4 |
| | Flakiness | F | 3 |
| • | Dryness | D | 2 |
| 15 | Wrinkling | W | 1 . |
| | Glazing | G | 1. |
| • | In order to obta | in a mum | 1 |

In order to obtain a numerical value for the total reaction at each site the severity score for each sign was multiplied by the rating. The resulting values were summed to give a total score for the degree of irritation at that site.

For example, the total score for a site assessed as R3, D2 and G1 would be:

$$(3 \times 3) + (2 \times 1) + (1 \times 1) = 12$$

The skin irritation

The skin irritation scores for the test and control products were compared statistically using the binomial probability test.

Immediately after assessment an identical fresh patch was applied to the same skin site for a further 24 hours (for a total of 48 hours) and skin reactions were again examined one hour after removal. The materials tested and the mean skin irritation scores after 24 hours and after 48 hours were as follows:

÷.

| | | | Skip lanians' |
|----|-------|---|-----------------|
| | Examp | le Material Tested | Skin Irritation |
| | 1 | Defonized water (control) | 24 hr. 48 hr. |
| | 2 | | 1.28 0.96 |
| - | _ | Trimethylglycine 50% w/v | 0.44 0.32 |
| 5 | • | in deionized water | |
| | 3 | sodium lauryl ether sulphate (SLES) 10.3% | 1.76 4.28 |
| | 4 | SLES 10.3% + trimethylglycine 7% | 0.56 0.80 |
| | 5 | SLES 10.3% + trimethylglycine 5% | 0.72 1.92 |
| | 6 | SLES 10.3% + trimethylglycine 3.5% | 0.76 2.40 |
| 10 | 7 | SLES 10.3% + trimethylglycine 2% | 1.12 2.92 |
| | 8 | SLES 5.15% + trimethylglycine 3.5% | 0.88 2.88 |
| | 9 | Deodorant (Control) | 0.96 |
| | 10 | Deodorant + 5% trimethylglycine | 0.65 |
| | 11 | Medical cleaner | 1.50 2.12 |
| 15 | 12 | Medical cleaner + 5% trimethylglycine | ·- <u>-</u> |
| | 13 | Eau de cologne | 1.27 1.50 |
| | 14 | | •• |
| | 15 | Esu de cologne + 5% trimethylglycine | •• |
| | | After Shave Lotion (Control) | 1.46 |
| 20 | 16 | After Shave Lotion + 5% trimethylglycine | 0.85 |

It can be seen from the foregoing numerical results that trimethylglycine significantly reduced the irritancy of the sample to the skin. Where a "--" is indicated in the chart above numerical results were not obtained.

Example 17

Transepidermal Water Loss (TEWL)

Irritants have much stronger effect on dry skin; thus ingredients of cosmetic compounds should not cause any increasing normal TEWL.

The effect of methanaminium, 1-carboxy-N,N,N-trimethyl-,hydroxide inner salt monohydrate on moisturizing and skin smoothness was tested by the tran-35 sepidermal water loss from skin (TEWL).

, ,

In example 17 a leading marketed skin cream was applied under an occlusive patch.

Skin Care Cream (commercially marketed)

| 5 | <u>Ingredients</u> | |
|----|---------------------------|---------|
| | Glycerol monostearate | 2.5 |
| | Cetyl stearyl ethoxylate | 5.0 |
| | Emulsifier (Stearic acid) | 2.0 |
| | White oil | 25.0 |
| 10 | Glycerol | 10.0 |
| , | Triethanolamine | 0.8 |
| | Water | Balance |

The same cream containing 50% w/w trimethylglycine was applied under an occlusive patch at a different skin site. As a control, blank occlusive patches were also applied to a third skin site. The transepidermal water loss from skin (TEWL) was measured.

The test subjects were twelve female volunteers

20 between 18 and 65 years of age. Panelists were selected
on the + criteria that they had normal healthy skin on
their forearms and were able to provide stable TEWL
values.

Three sites were marked on the volar forearm of
each volunteer at approximately equal intervals between
the elbow crease and the wrist. The subject was then
seated comfortably with her forearms exposed for a period
of half an hour so that they would equilibrate with
conditions of temperature and relative humidity in the
laboratory.

TEWL readings were taken from each skin site to provide normal untreated water loss levels. Treatment was made via 12 mm occlusive Finn chambers. A blank Finn chamber served as the control site. The lotion samples were applied by immersing Whatman 3MM filter paper discs in the sample, removing excess product against the edge

of the container and placing the disc in the Finn chamber using forceps. Each chamber was applied to one of the marked sites on the forearm and held in position with scanpor non-occulsive surgical tape. The pattern of application of treatments to sites was randomized and recorded.

18 to 24 hours later each subject returned and was re-equilibrated as above. The chambers were then removed and the skin was blotted with tissue to absorb

10 surface moisture. TEWL readings were recorded at 1 minute intervals from the time of removal of treatment until a steady valve was obtained, a period of ten to twenty-five minutes depending on the individual concerned and the treatment applied.

- Temperature and relative humidity were recorded at the start of each TEWL reading as gross changes can significantly alter the rate of water loss. The results for each panelist are presented graphically below. An analysis of the results revealed the following:
- 1) The 1 minute TEWL reading immediately after patch removal is lower for the cream containing trimethylglycine than for the cream alone indicating that the trimethylglycine-treated sites lost water less rapidly. The blank patch control is lower still and represents the skin hydrating effect of occlusion, without adding additional moisture to the skin.
- 2) The final TEWL value obtained when readings approach a constant value is lower for the cream containing trimethylglycine than for the cream alone, and frequently also the blank control, again indicating better moisture retention at trimethylglycine-treated skin sites.
- 3) The rate at which the TEWL value decreases towards the normal untreated level is in general more rapid for the cream containing trimethylglycine than for the cream alone or the blank control.

These results are summarized in the table below:

| | | Ir | nitial TEWL leve (g/m ² h) | | evel | | Final TEWL level (g/m h) | | | | | | Time for 75 % reduction in TEWL rate (min) | | | |
|----|----|------------|--|------|----------|------------|--------------------------|----|-----|----|-----|------|--|----------------|--------------|--|
| | 5 | Par No. | _ | lenk | Cre | 8 <i>m</i> | Cream +TMG | 8 | ank | Cr | eam | Cres | | k Crea | m Cream | |
| | | 1 | `. • | | 58 : | | 40 | 7 | | 8 | | 5 | * | | | |
| • | LO | 2 | 36 | | 50 | • | 48 | 10 | | 12 | | 8 | Not | 7 14 | 5 7 | |
| | | 3 | 34 | | | | | | | | | | reach | ed | , | |
| | | 4 | 20 | | 41 41 | | i 4 | 8 | | 9 | | 7 | 16 | 12 | 6 | |
| | | | | • | •1 | | | 7 | | 12 | | 8 | Not | Not | • | |
| | 5 | ; | 30 | 6 | 3 | | | | | | | | reache | d reache | d | |
| 1. | 5 | | | | • | 39 | • | 8 | | 18 | | 8 | Not | Not | 13 | |
| | 6 | | 17 | 6 | 1 | 53 | | 9 | | | | | reached | reached | 1 | |
| | | | | | | | | , | | 20 | 8 | В | Not | 13 | 8 | |
| | 7 | | 17 | 78 | ; | 45 | | 7 | | 22 | 3 | | reached | | | |
| | | | | | | | | | | _ | J | , | Not | Not | - | |
| 20 | 8 | | 26 | 71 | | 65 | • | 11 | 2 | 6 | 1: | 1 | reached Not | · Gaerieg | | |
| | _ | | | | | • | | | | | | | | Not reached | 7 | |
| | 9 | | 25 | 74 | | 63 | 6 | | 19 | 9 | 8 | | 11 | Not | - | |
| | 10 | | 18. | | | | | | | | | | | reached | 7 | |
| 5 | | | . 10 | i | | 34 | 7 | | i | | 6 | | Not | i | 7 | |
| - | 11 | | 23 | 53 | | | | | | | | | reached | | • | |
| | | | | -3 | | 53 | 10 | | 17 | | 9 | ł | Not | Not | 8 | |
| | 12 | | 26 | 54 | , | 30 | | | | | | ı | reached | reached | | |
| | | | | -7 | | 30 | 6 | | 13 | | 4 | 1 | 17 | 18 | 7 | |

The "*" indicates that the evapormeter probe shifted over in the early part of curve so readings were not valid.

35 The "i" indicates that the patch came loose over the application period.

· .

q.s. to pH 6.5-7.2

It can be seen that under these test conditions there were clear indications that the addition of trimethylglycine to the skin cream resulted in improved moisture retention when the product was applied to the skin.

5

Example 18

Example 18 provides illustrative formulations including ingredient with irritating properties for a skin cream, body shampoo and skin cleanser in accordance with the invention.

Skin cream

| Ingredient | |
|--|------------|
| Trioleyl phosphate | Amount (%) |
| 15 Petrolatum | 3.0 |
| Glyceryl stearate | 18.0 |
| Isopropyl palmitate | 5.0 |
| Cetyl alcohol | 4.0 |
| Stearyl heptanoate (irr.) | 2.0 |
| 20 Cetearyl catalante (1rr.) | 0.5 |
| 20 Cetearyl octanoate (irr.) Sorbitol | 0.5 |
| | 5.0 |
| Trimethylglycine | 5.0 |
| Water, perfume, preservative (irr.) | q.s. |
| 25 Shampoo | |
| Ingredient | |
| · | Amount (%) |
| Sodium lauryl ether sulphate 30% (irr.) Coconut monoethanolamide | 40.0 |
| Trimethylglycine | 2.0 |
| 30 Perfume, colour, water | 5.0 |
| HC1/NaOH | q.s. |
| · · · · · · · · · · · · · · · · · · · | |

Sodium lauryl ether sulphate is known to have a hard 35 cleansing effect; thus because of the side effect (e.g. too effective fat removal from the skin and hair) it

:

cannot be used as such in shampoos. The addition of trimethylglycine remarkably reduces the irritation effect.

Skin cleanser for oily skin

| _ | Tor olly skin | |
|-----|-------------------------------------|------------|
| 5 | Ingredient | - |
| | Triclosan (deodorising agent; irr.) | Amount (%) |
| 10 | (IFF.) | 3.0 |
| | DEA-oleth-3 phosphate | 10.0 |
| | Hydroxypropylcellulose | 2.5 |
| | Amphoteric-1 | 2.5 |
| | Trimethylglycine | 5.0 |
| | Water | 5.0 |
| | Ethanol (96%) | 37.0 |
| _ | | 35.0 |
| 7 = | | |

Trimethyl glycine reduces the irritating properties of irritants.

Example 19 - Smoothness Test

Body Lotion With Improved Skin Compatibility

| 20 | Ingredient | <u>ibility</u> |
|-----|-------------------------------|----------------|
| | Emulsifier | Amount (%) |
| 25 | Cetylalcohol (irr.) | 2.0 |
| | Isopropylstearate | 2.5 |
| | Preservative (potential irr.) | 4.5 |
| | ugret | 0.03 |
| | 1,2-Propyleneglycol | 78.77 |
| | Trimethylglycine | 6.0 |
| | Parfum oil | 0 or 6.0 |
| 30 | | 0.2 |
| 311 | | |

The effect of methanaminium, 1-carboxyN,N,N-trimethyl-,hydroxide inner salt monohydrate on skin
tested by the Zeiss-spektrophotometer. This formula was
tested by using ten female volunteers. The body lotion
was used for 8 days. The smoothness of the skin was
measured by a Zeiss-sp ktrophotometer by 660 nm. The
results show that the addition of trimethylglycine moderately improves the smoothness of the skin.

Smoothness of skin was calculated by the following equation:

= <u>extinction after treatment</u> initial extinction

X 100

5 Results for the ten subjects are reported below:

| | | |
|----------------------|-------|---------------------|
| Age of the volunteer | 0% | 6% Trimethylglycine |
| 43 | 99.8 | 01.0 |
| 44 | 99.3 | 91.8 |
| 51 | | 89.4 |
| 50 | 75.9 | 59.4 |
| | 98.6 | 79.3 |
| 70 | 97.5 | 89.4 |
| 47 | 98.3 | 91.9 |
| 34 | 100.3 | |
| 64 | | 97.1 |
| 30 | 75.4 | 97.8 |
| 51 | 100.0 | 97.4 |
| <u> </u> | 95.3 | 96.5 |
| Average | 94.0 | 89.0 |
| | | |

Example 20 - Moisture Holding Capacity Using Corneometer

The same body lotion formula used in Example 19
was used in tests where the moisturizing effect of trimethylglycine was measured by a Corneometer. The moisture
holding capacity in the skin was increased when the
trimethylglycine level in the product increased from 0% to
2% to 6%.

The same group of test persons and model products

were used as in Example 19. Medial part of skin of arms

were used (5 x 2.5 cm²). Different test areas for different model products were used. The instrument used for

measurement of the lotions, skin moisturizing effect was a

Corneometer Combi (Courage & Khazaka Electronic GmbH,

FRG). The subjects were acclimatized in a controlled

atmosphere (vH 60 %, T 24 °C) for 30 minutes before 0-values of different skin areas were measured. Body lotions were used twice daily (12 hr. intervals). Duration of the trials was 8 days.

5 The following are hydration measurements made by the Corneometer Combi:

| | Product | Measu | rement | | | | | | | |
|----|--|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|----------------------------------|
| 10 | | | after Appl | 10 | 20 | 30 | Minut 40 | es 60 | 90 | 120 |
| 15 | Control 0 % 2 % 6 % trimethyl- slycine | 107.8 104.5 104.0 103.6 | 106.3 155.8 152.9 150.4 | 106.8 118.5 120.0 122.1 | 107.8 118.4 119.2 118.7 | 107.7 116.1 115.9 116.1 | 108.7 115.7 117.1 117.6 | 107.6 113.2 113.8 116.1 | 108.2 112.6 113.8 116.3 | 107.8 111.8 112.8 114.9 |

20

25

30

PATENT CLAIMS:

1. A method of reducing the skin irritating properties of a cosmetic composition containing substantially no hydrolyzed protein, characterized 5 by the methanaminium, 1-carboxy-N, N, N-trimethyl-, hydroxide, inner salt monohydrate having the chemical struc-

or an anhydride thereof is incorporated into the cosmetic composition in an amount effective to neutralize the skin-irritating properties of the cosmetic composition.

- 2. The method according to claim 1 20 characterized by that the methanaminium, 1-carboxy-N, N, N-trimethyl-, hydroxide, inner salt is added in concentration of 0,1 - 50%, preferably 0,1 - 15%, percent by weight.
- 3. A cosmetic composition having improved skin 25 compatibility and moisturizing properties containing substantially no hydrolysed protein and character i z e d by that it comprises a compound having irritating or sensitizing properties when applied to human skin, 30
- an effective amount of a quaternary ammonium like compound, methanaminium, 1-carboxy-N, N, N-trimethyl-, hydroxide, inner salt monohydrate having the chemical

- 5 or anhydride thereof.
 - .4. The cosmetic composition of claim 3 or 4 characterized by that it is a skin cleanser, a skin cream, or body shampoo.
- The cosmetic composition of claim 3 or 4 characterized by that the said compound is a 10 deodorant.

The cosmetic composition of claim 3, wherein the concentration of methanaminium, 1-carboxy-N,N,N-trimethyl-, hydroxide, inner salt monohydrate is from 0.1 to 50,

preferably from about 0.1 to 15 percent by weight. 15

The cosmetic composition of claim 3 or 4 characterized by that it is a cream, ointment, emulsion, or an aerosol.

- 6. A topical preparation according to claim 3 and having improved skin compatibility and moisturizing 20 properties characterized by that it com-
 - (a) one or more cosmetic ingredients selected from the group consisting of an active agent, an
- emollient, a barrier agent, a healing agent, a humectant, 25 an emulsifier, a preservative, a perfume oil, a coloring agent, a deodorising agent, a surfactant, a solvent and mixtures of the above; and
- (b) one or more cosmetic ingredients [may be included in group (a) and] having irritating properties 30 when applied to human skin; and
 - (c) trimethylglycine or anhydride thereof in an amount effective to neutralize the skin-irritating effects of said cosmetic ingredient.

1.

INTERNATIONAL SEARCH REPORT

| 1 | I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all PC5: A 61 K 7/40 7/67 | PCT/ET 01/00175 |
|--------------|--|--|
| - 1 | According to International Patent Classification (IPC) or to both National Classification and IPC IPC5: A 61 K 7/48, 7/075, 7/50, 7/00 | D ₆ |
| 1. | IPC5: A 61 K 7/48, 7/075, 7/50, 7/00 | ., |
| | II. FIELDS SEARCHED | |
| · L | | |
| <u> </u> | lassification System Minimum Documentation Searched 7 | |
| - 1 | Classification Symbols | |
| 171 | PC5 | |
| - | A 61 K | |
| - 1 | Documentation Co. | |
| | . Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in Fields Searched ⁸ | |
| - 1 | | |
| SE | ,DK,FI,NO classes as above | |
| 111. | DOCUMENTS CONSIDERED TO BE RELEVANTS | |
| Cate | gory • Citation of Decument 11 and 12 | |
| X | WO, A1, 8202337 (WELLA AKTIENCEST), of the relevant passages 12 | In |
| - 1 | WO, A1, 8202337 (WELLA AKTIENGESELLSCHAFT) 22 July 1982, | Relevant to Claim No.13 |
| - 1 | see the whole document | 1-6 |
| - 1 | ord document | |
| x | DE A1 PERSON | 1 |
| | DE, A1, 3527974 (WELLA AG) 12 February 1987, | |
| - 1 | see the whole document | 1-6 |
| lx | | 1 |
| ^ | DE, A, 1911144 (MEDISAN AB) 17 September 1970, see the whole document | |
| - 1 | see the whole document | 1-6 |
| 1 | | - 0 |
| X | Patent Abstracts . C . | |
| 1 | Patent Abstracts of Japan, Vol 7, No 191, C182, abstract of JP 58- 92607, publ 1983-05-06 | |
| 1 | abstract of JP 58- 92607, publ 1983-06-02 | 1-6 |
| x | · | |
| 1 | FR, A, 2303536 (RHONE-POULENC INDUSTRIES) 8 October 1976. | |
| 1 | 8 October 1976, see the whole document | 1-6 |
| 1 . | ano le document | |
| 1 | | 1 |
| · | | |
| N Committee | | |
| "A" doc | al categories of cited documents: 10 | |
| con: | sidered to be of narrial state of the art which | iternational files |
| | of or after the internation invention | to the application but heory underlying the |
| whia | Comment of socialists | claimed invention |
| "O" docu | on or other special reason (as specified) ment referring to a specified on the publication date of another special reason (as specified) cannot be considered novel or cannot be conside | pe considered to |
| *P* docur | media tradition the section of Continored to Tradition of Continored to Tra | claimed invention |
| /. CERTIFI | ment published prior to the international filing date but CATION document is combined with one or ments, such combination being obvious document is combined with one or mount in the arm document is combined with one or mount in the arm document is combined with one or mount in the arm document is combined with one or mount in the arm document is combined with one or mount in the arm document is combined with one or mount in the arm of the | re other such docu- is to a person skilled |
| ata of the A | CATION Category Completion of the International Search Category Category Completion of the International Search Category Categ | family |
| 2th Sen | THE HILL HALL COMMENT | |
| | | eport . |
| ernational S | earching Authority | 1 " |
| | Signature of Authorized Officer | |
| PCT/ISA/62 | SWEDISH PATENT OFFICE (second sheet) (January 1985) Elisabeth Carlbons | > |
| | (second sheet) (January 1985) Elisabeth Carlborg | |

| III. DOCU | MENTS CONSIDERED TO BE RELEVANT Citation of Document | - International | Application No. DCT | /FI 91/00176 |
|--------------------|---|------------------|---------------------|----------------------|
| Category - | Citation of Des | (CONTINUES - | THO PU | /FI 91/00176 |
| 1 1 | with indication | D. When a | THE SECOND SHEET | 1 |
| x | EP, A2, 0186453 (WARNER-L 2 July 1986, see the whole document | of the | elevant passages | |
| - 1 - 1 | 2 .1/11 100453 (WARNER-L | AMBERT COVOLUM | | Relevant to Claim No |
| - 1 - 1 | 200 +1-1986, | COMPANY) | | |
| 1 1 | see the whole document | t | | 1-6 |
| - 1. 1 | | | | 1 |
| X | P, A2, 0047916 (TH. GOLDS 24 March 1982, | ~ ~ | | 1 |
| 1 1 | 24, 0047916 (TH. GOLDS | CLINTE | | |
| | 24 March 1982, | CUMITAL AG) | | |
| 1 1 | see the whole document | | | 1-6 |
| 1. | | | - 1 | |
| X DE | , AI, 2750731 (REWO CHEMI 17 May 1979, see the whole | | 1 | |
| | , 71, 2/50731 (REWO CHEMI | SCUP II | - 1 | |
| | 17 may 1979, | SCHE WERKE GMBH) | | |
| 1 1 | see the whole document | • • • | 1 | 1-6 |
| 1 1 | | _ | · | |
| | | | - 1 | i |
| 1 1 | - | · | 1 | |
| i 1 | | | 1 | 1 |
| 1 1 | | | 1 | - 1 |
| 1 1 | | | 1 | I |
| 1 1 | • | | 1 | 1 |
| 1 1 | | | . | |
| - | | | 1 | |
| | | | 1. | |
| | | | | |
| - 1 | | | 1 | 1 |
| | | | 1 | |
| 1 | | | 1. | 1 |
| 1 | | | 1 | 1 |
| 1 | | | | 1 |
| · [. | | | | |
| 1 | | | . | 1 |
| | | | | |
| 1 | | | 1 | 1 |
| 1 | | | 1 | 1 |
| 1 | | • | | 1 |
| - 1 | | | 1 | 1 |
| 1 | | | 1 | 1 |
| - 1 | | | 1 | 1 |
| - 1 | • | • | 1 | 1 |
| - 1 | • | | 1 | 1 |
| - 1 | | | 1 | 1 |
| | | • | 1 | 1 |
| 1 | | • | 1 | . 1 |
| 1 | | | 1 | 1 |
| 1 | | | 1 | 1 |
| 1 | | • | 1 | 1 |
| 1 | | | 1 | 1 |
| | | | 1 | 1 |
| 1 | | | - 1 | 1 |
| | • | | 1 | 1 |
| | | | | i |
| | | | | 1 |
| | • | | | |
| | | | | |
| | | • | | |
| | | • . | | |
| W219 (over | | • . | | |
| W/210 (extra shee) | 3 (January 1985) | • | | |
| A/210 (extra sheet | (January 1985) | | | |

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/FI 91/00176

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the Swedish Patent Office EDP file on 91-07-31

The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

| WO-A1- 8202337 | | | Patent family member(s) | Publication |
|----------------|----------|---|--|--|
| | 82-07-22 | AT-E- AU-B- | 10063 550795 | 84-11-15 |
| | | AU-D- DE-A-C EP-A-B US-A- | 8001582 3101011 0056595 | 86-04-10 82-08-02 82-07-22 82-07-28 |
| DE-A1- 3527974 | 87-02-12 | NONE | 4752467. | 88-06-21 |
|)E-A- 1911144 | 70-09-17 | AT-A- | 294319 | 71-10-15 |
| | | AT-A-B- BE-A- CH-A- CH-A- | 306928 729465 513648 576789 | 73-03-15 69-08-18 71-10-15 |
| | | FR-A- GB-A- LU-A- | 2003312 1232569 58143 | 76-06-30 69-11-07 71-05-19 |
| | | NL-A- US-A- | 6903383 3666863 | 69-07-07 69-09-09 72-05-30 |
| -A- 2303536 | 76-10-08 | BE-A- CH-A- DE-A- GB-A- JP-A- LU-A- NL-A- | 839402 594604 2610225 1479132 51113829 74518 7602210 | 76-09-10 76-09-10 78-01-13 76-09-23 77-07-06 76-10-07 77-01-11 76-09-14 |
| A2- 0186453 | 86-07-02 | US-A- AU-D- | 4058597 5078885 | 77-11-15 |
| 2- 0047916 | | JP-A- US-A- | 61171800 4780249 | 86-08-02 88-10-25 |
| l- 2750731 | 82-03-24 | DE-A-C- US-A- | 3033929 4420410 | 82-03-18 83-12-13 |
| | 79-05-17 | GB-A-B- US-A- | 2010832 4221733 | 79-07-04 80-09-09 |